

Production of 2-phenylethanol (PEA) by yeast with ionic liquids in situ extraction

Patrycja OKUNIEWSKA*, Urszula DOMAŃSKA-ŻELAZNA, Aneta POBUDKOWSKA – Department of Physical Chemistry, Faculty of Chemistry, Warsaw University of Technology, Warsaw, Poland; Jolanta MIERZEJEWSKA – Department of Drug Technology and Biotechnology, Institute of Biotechnology, Faculty of Chemistry, Warsaw University of Technology, Warsaw, Poland

Please cite as: CHEMIK 2016, 70, 9, 491–496

Introduction

PEA is a valuable aroma of a subtle rose scent used in food, perfume and household chemicals, as well as a preservative (in medicines) and a disinfectant (e.g. for strawberries). High demand for PEA motivates the ongoing quest for new production methods. An alternative for synthetic PEA is obtaining this fragrance from rose petals. However, the costs of such production are very high – the price of rose oil (containing approx. 66–79% PEA [1]) is approx. 4,600 euro/kg [2]. The paper presents the application of yeasts for the PEA synthesis. The product of such synthesis, under the applicable law, is considered as a natural product [3]. The only problem is the low final concentration of the product. This can be solved using the *in situ* extraction [4]. There are the following methods for the PEA *in situ* extraction available: adsorption [5–8], microcapsules [9, 10], membrane extraction [11], liquid-liquid extraction using oleic acid [12, 13] or ILs [4, 9, 14–17]. These methods allow for the increase of the production of PEA to approx. 6 g/L in case of flask cultures and 12 g/L to 26 g/L for fed-batch cultures [18]. The study focuses on the liquid-liquid extraction using ionic liquids due to the possibility to match the ionic liquid for the purposes of research and the simplicity of this solution. The possibility to re-use the ionic liquid after the PEA re-extraction was another motive. The aim of the studies was to find the best ionic liquid for the extraction of PEA *in situ*.

Description

Experimental part

Materials

The studies involved use of YPD broth for yeast growth (20 g/L glucose, 10 g/L yeast extract, 20 g/L peptone), while the experiment itself was conducted using production broth (10 g/L glucose, 10 g/L sucrose, 6 g/L L-phenylalanine (L-Phe), 4 g/L KH_2PO_4 , 0.4 g/L $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$, 1 g/L yeast extract). The studies involved the strain *Saccharomyces cerevisiae* AM1 d from the collection of the Department of Drug Technology and Biotechnology. The following *in situ* extractants were used [C_8jQuin][NTf_2] (the synthesis method and purity were presented in the paper [19]) [BMPyr][NTf_2] (Io-li-tec, >0.99), [$\text{P}_{6,6,6,14}$][TCM] (MERCK KGaA, ≥0.98 [20]), [HMPIP][NTf_2] (Io-li-tec, >0.99*).

Methods

3 yeast colonies were transferred into a 100 mL Erlenmeyer flask containing 10 mL of YPD broth. The culture was left in an incubating shaker (30°C, 220 rpm, Lab Companion SI-600R). On the next day, the over-night culture was used to inoculate production broth in such a manner so the initial optical density of culture is 0.2,

when measured at wavelength $\lambda=600$ nm (OD_{600}). The culture was cultivated in the incubating shaker (30°C, 220 rpm). OD_{600} of culture was measured every 1 hour. When OD_{600} reached 1, the culture was poured into 4 flasks (100 mL): 15 mL to each reference flask and 12 mL to each test flask, followed by the addition of 3 mL of the extractant to the test flasks. The cultures were cultivated in two biological replicates. In the extractant-free sample, the yeast growth was measured (OD_{600} measurements for 8 hours). The cultures were centrifuged (25°C, 1000 g, 2 min, Eppendorf Centrifuge 5804R) after 24 hours and 48 hours in order to separate the phases and OD_{600} of the aqueous phase measurement. Then, they were re-centrifuged (25°C, 2000 g, 3 min) in order to separate biomass. The phases were collected into test tubes, followed by the addition of an equal volume of acetonitrile (2-fold dilution). The final concentration of 2-phenylethanol in both phases was determined by means of HPLC (Agilent Technologies 1200 series, column C18, 25°C, 1 mL, acetonitrile:water = 50:50) at wavelength $\lambda=220$ nm.

Result discussion

Completely new ionic liquids were used as potential PEA *in situ* extractants. The results of cultures presented (Fig. 1) clearly show that the addition of ionic liquids may have various effects on yeast growth. Two ionic liquids [BMPyr][NTf_2] ($\text{OD}_{600}^{24\text{h}}=1.78$, $\text{OD}_{600}^{48\text{h}}=1.92$), and [HMPIP][NTf_2] ($\text{OD}_{600}^{24\text{h}}=1.78$, $\text{OD}_{600}^{48\text{h}}=1.86$) intensified the yeast growth, while the other two [C_8jQuin][NTf_2] ($\text{OD}_{600}^{24\text{h}}=1.36$, $\text{OD}_{600}^{48\text{h}}=1.50$) and [$\text{P}_{6,6,6,14}$][TCM] ($\text{OD}_{600}^{24\text{h}}=1.42$, $\text{OD}_{600}^{48\text{h}}=1.43$) slowed down the yeast growth in comparison with the reference ($\text{OD}_{600}^{24\text{h}}=1.64$, $\text{OD}_{600}^{48\text{h}}=1.65$), as it is clearly shown in Figure 1. The slowed growth could have been caused by a need of additional adaption of yeast to these specific ionic liquids. On the other hand, the intensified growth is probably a result of low PEA concentration in the aqueous phase, thus eliminating its inhibiting effect on the yeast growth occurring for the concentrations of approx. 2–3 g/L [21].




Fig. 1. Culture growth presented by optical density at wavelength $\lambda = 600\text{nm}$

*Corresponding author:

Patrycja OKUNIEWSKA – M.Sc., (Eng.), e-mail: pokuniewska@ch.pw.edu.pl

For three of the ionic liquids tested after 24h of the cultivation, the PEA concentrations obtained (in grams per litre of culture) were much higher than for the reference sample (no extractant) – 2.10 ± 0.009 g/L. These were in the descending order [BMPyr][NTf₂] (in the IL phase 6.71 ± 0.006 g PEA/L), [HMPIP][NTf₂] (in the IL phase 6.21 ± 0.010 g/L), [C₈iQuin][NTf₂] (in the IL phase 3.89 ± 0.004 g/L). For [P_{6,6,6,14}][TCM], the PEA concentration in the IL phase after 48-hour culture was only 0.05 ± 0.005 g/L. 2-phenylethanol was still produced, which is indicated by much higher concentrations after 48 hours. The PEA concentration in samples after 48 hours depending on the ionic liquid in the descending order: [BMPyr][NTf₂] (in the IL phase 10.25 ± 0.019 g PEA/L), [HMPIP][NTf₂] (in the IL phase 6.21 ± 0.009 g/L), [C₈iQuin][NTf₂] (in the IL phase 3.89 ± 0.007 g/L), reference (no extractant) 2.10 ± 0.014 g/L, [P_{6,6,6,14}][TCM] (in the IL phase 1.92 ± 0.007 g/L). The results are presented in Figure 2.




Fig. 2. PEA concentration after 24 hours and 48 hours in grams per litre of culture in the aqueous phase (a.p.) and the ionic liquid phase (IL phase)

The bioconversion results L-Phe to PEA (Fig. 3) at 70% for [BMPyr][NTf₂] after 48 hours show that a further conversion was still possible. Therefore, we plan to set new 72-hour cultures. The organic phase/aqueous phase partition coefficients for PEA (Fig. 4) indicate a good PEA extraction by the selected ionic liquids zamiast good PEA extraction by the ionic liquids selected. For [P_{6,6,6,14}][TCM] after 48 hours, the value of the partition coefficient was at its maximum, however, as already mentioned, the total PEA concentrations in the culture were very low, which makes it useless for the *in situ* extraction.




Fig. 3. Bioconversion percentage: L-phenylalanine (L-Phe) to 2-phenylethanol (PEA)




Fig. 4. IL phase/aqueous phase partition coefficient (K_D) for 2-phenylethanol (PEA)

Summary and conclusions

None of the ionic liquids studied proved to be toxic for the yeast *Saccharomyces cerevisiae* AM1d. The results presented clearly show that the best extractant for the *in situ* extraction of PEA is butyl-methylpyrrolidinium bis(trifluoromethylsulfonyl)imide [BMPyr][NTf₂]. This is shown by a high L-Phe conversion (50% after 24 hours and 70% after 48 hours), high PEA concentration in the ionic liquid phase (6.71 ± 0.006 g/L after 24 hours and 10.25 ± 0.019 g/L after 48 hours), as well as a high IL phase/aqueous phase partition coefficient for PEA ($K_D = 12.22$ after 24 hours and $K_D = 19.4$ after 48 hours). Another promising extractant is [HMPIP][NTf₂], as indicated by a high L-Phe conversion (43% after 24 hours and 35% after 48 hours), close to the reference sample, high PEA concentration in the ionic liquid phase (4.74 ± 0.009 g/L after 24 h and 6.21 ± 0.010 g/L after 48 h), as well as a high IL phase/aqueous phase partition coefficient for PEA ($K_D = 17.01$ after 24 h and $K_D = 12.97$ after 48 h). However, the conversion percentage of L-Phe (a maximum of 70% after 48 hours) suggests that the culture should be cultivated for longer (e.g. for 72 hours) in order to fully use the substrate. The ionic liquids studied in this work proved to be much better than previously tested ionic liquids containing the same anion and the following cations: 1-benzyl-3-methylimidazolium (~ 1.2 g PEA/L, $K_D = 17.6$), 1-methyl-1-propylpiperidinium (~ 1.2 g PEA/L, $K_D = 11.6$), methyltriethylammonium (~ 1.0 g PEA/L, $K_D = 4.0$) [15].

Acknowledgements

This work has been supported by the project of National Science Center in Poland for years 2015–2018, No. 201/15/B/ST5/00136

Literature

- Verma R. S., Padalia R. C., Chauhan A., Singh A., Yadav A. K.: *Volatiles Constituents of Essential Oil and Rose Water of Damask Rose, Rosa Damascena Mill., Cultivars from North Indian Hills*. Nat. Prod. Res. 2011, **25**, 17, 1577–1584.
- Kovacheva N., Rusanov K., Atanassov I.: *Industrial Cultivation of Oil Bearing Rose and Rose Oil Production in Bulgaria During 21ST Century, Directions and Challenges*. Biotechnol. Equip. 2010, **24**, 2, 1793–1798.
- Directive of the European Parliament and of the Council No. 1334/2008/EC December 16, 2008.
- Okuniewska P., Ramjugernath D., Naidoo P., Domańska U.: *Solubility of Ionic Liquids in 2-Phenylethanol PEA. and Water*. Fluid Phase Equilibria 2014, **376**, 55–63.

5. Carpiné D., Dagostin J. L. A., da Silva V. R., Igarashi-Mafra L., Mafra M. R.: *Adsorption of Volatile Aroma Compound 2-Phenylethanol from Synthetic Solution onto Granular Activated Carbon in Batch and Continuous Modes*. *J. Food Eng.* 2013, **117**, 3, 370–377.
6. Šimko I., Roriz E., Gramblička M., Illeová V., Polakovič M.: *Adsorption Separation of 2-Phenylethanol and L-Phenylalanine on Polymeric Resins: Adsorbent Screening, Single-Component and Binary Equilibria*. *Food Bioprod. Process.* 2015, **95**, 254–263.
7. Mei J.: *Enhanced Biotransformation of L-Phenylalanine to 2-Phenylethanol Using an in Situ Product Adsorption Technique*. *Process Biochem.* 2009, **44**, 8, 886–890.
8. Hua D., Lin S., Li Y., Chen H., Zhang Z., Du Y., Zhang X., Xu P.: *Enhanced 2-Phenylethanol Production from L-Phenylalanine via in Situ Product Adsorption*. *Biocatal. Biotransformation* 2010, **28**, 4, 259–266.
9. Stark D., Kornmann H., Munch T., Sonnleitner B., Marison I. W., von Stockar U.: *Novel Type of in Situ Extraction: Use of Solvent Containing Microcapsules for the Bioconversion of 2-Phenylethanol from L-Phenylalanine by Saccharomyces Cerevisiae*. *Biotechnol. Bioeng.* 2003, **83**, 4, 376–385.
10. Gao F., Daugulis A. J.: *Bioproduction of the Aroma Compound 2-Phenylethanol in a Solid-Liquid Two-Phase Partitioning Bioreactor System by Kluyveromyces Marxianus*. *Biotechnol. Bioeng.* 2009, **104**, 2, 332–339.
11. Mihal' M., Vereš R., Markoš J.: *Investigation of 2-Phenylethanol Production in Fed-Batch Hybrid Bioreactor: Membrane Extraction and Microfiltration*. *Sep. Purif. Technol.* 2012, **95**, 126–135.
12. Stark D., Münch T., Sonnleitner B., Marison I. W., von Stockar U.: *Extractive Bioconversion of 2-Phenylethanol from L-Phenylalanine by Saccharomyces Cerevisiae*. *Biotechnol. Prog.* 2002, **18**, 3, 514–523.
13. Etschmann M. M. W., Sell D., Schrader J.: *Screening of Yeasts for the Production of the Aroma Compound 2-Phenylethanol in a Molasses-Based Medium*. *Biotechnol. Lett.* 2003, **25**, 7, 531–536.
14. Wang H., Dong Q., Meng C., Shi X. A., Guo Y.: *A Continuous and Adsorptive Bioprocess for Efficient Production of the Natural Aroma Chemical 2-Phenylethanol with Yeast*. *Enzyme Microb. Technol.* 2011, **48**, 4–5, 404–407.
15. Sendovski M., Nir N., Fishman A.: *Bioproduction of 2-Phenylethanol in a Biphasic Ionic Liquid Aqueous System*. *J. Agric. Food Chem.* 2010, **58**, 4, 2260–2265.
16. Xiang Z. Y., Lu Y. C., Zou Y., Gong X. C., Luo G. S.: *Preparation of Microcapsules Containing Ionic Liquids with a New Solvent Extraction System*. *React. Funct. Polym.* 2008, **68**, 8, 1260–1265.
17. Domańska U., Zawadzki M., Królikowski M., Lewandowska A.: *Phase Equilibria Study of Binary and Ternary Mixtures of {N-Octylisoquinolinium Bis{trifluoromethyl.sulfonyl}imide + hydrocarbon, or an Alcohol, or Water}*. *Chem. Eng. J.* 2012, **181–182**, 63–71.
18. Hua D., Xu P.: *Recent Advances in Biotechnological Production of 2-Phenylethanol*. *Biotechnol. Adv.* 2011, **29**, 6, 654–660.
19. Domańska U., Zawadzki M., Królikowska M., Marc Tshibangu M., Ramjugernath D., Letcher T. M.: *Measurements of Activity Coefficients at Infinite Dilution of Organic Compounds and Water in Isoquinolinium-Based Ionic Liquid [C₈iQuin][NTf₂] Using GLC*. *J. Chem. Thermodyn.* 2011, **43**, 3, 499–504.
20. Domańska U., Okuniewska P., Markowska A.: *Phase Equilibrium in Binary Systems of Ionic Liquids, or Deep Eutectic Solvents with 2-Phenylethanol PEA, or Water*. *Fluid Phase Equilibria* 2016, **424**, 68–78.
21. Stark D., Zala D., Münch T., Sonnleitner B., Marison I. W., von Stockar U.: *Inhibition Aspects of the Bioconversion of L-Phenylalanine to 2-Phenylethanol by Saccharomyces Cerevisiae*. *Enzyme Microb. Technol.* 2003, **32**, 2, 212–223.

Patrycja OKUNIEWSKA – M.Sc., (Eng.), (M.Sc. in 2012 at the Faculty of Chemistry/Industrial Biotechnology, Warsaw University of Technology).
 Doctorate at the Faculty of Chemistry at Warsaw University of Technology (2012 – ongoing). Number of publications: 5 Research interests: Extraction, cultures of yeast, 2-phenylethanol.

e-mail: pokuniewska@ch.pw.edu.pl, phone: +48698692767

Professor Urszula DOMAŃSKA-ŻELAZNA – Ph.D., D.Sc., (Eng.), (Faculty of Chemistry, Warsaw University of Technology). Since 1968 works in the Department of Physical Chemistry, Faculty of Chemistry, Warsaw University of Technology. From 1991 to 1995, and since 2000 until now Head of the Department of Physical Chemistry. Teaches physical chemistry and thermodynamics of phase equilibria. Awards/prizes: 2 Minister of Education, Scientific Award of Warsaw University of Technology, 25 Awards of the Rector of the Warsaw University of Technology. Number of publications: 350 in foreign magazines and 5 monographs or chapters in scientific books. Research interests: thermodynamics of phase equilibria, solid-liquid equilibria, calorimetry, correlation and prediction of physicochemical properties, ionic liquids, pharmaceuticals and fragrances.

e-mail: ula@ch.pw.edu.pl, phone: +48(22)6282741

Aneta POBUDKOWSKA – Ph.D., (Eng.), (M.Sc. in 2001, Faculty of Chemical and Process Engineering, Warsaw University of Technology, Ph.D. in chemistry in 2006, Faculty of Chemistry, Warsaw University of Technology). In the years 2004–2010 worked as a technical in the Department of Physical Chemistry of the Faculty, and from 2010 – appointed Assistant Professor in the same team. Teaches physical chemistry and physical chemistry of drugs. Publications: 27 scientific publications in foreign journals. Awards/ prizes: 3 times Individual Award of Rector of Warsaw University of Technology, Scholarship of Ministry of Science and Higher Education for Outstanding Young Scientists and Scholarship Conference for Young Workers Scientific Research awarded by the Society for Scientific Warsaw and the Foundation for Polish Science, 2 times Golden Chalk Award, Medal of the Memorial for special merits for the development of teaching, shaping the image and promotion of the Faculty of Chemistry of Warsaw University of Technology. Research interests: solid-liquid equilibria and physicochemical properties of pharmaceuticals.

e-mail: pobudka@ch.pw.edu.pl, phone: +48(22)234 7475

Jolanta MIERZEJEWSKA – Ph.D., (M.Sc. in biological sciences, specializing in microbiology, 2002., University of Warsaw; Ph.D. in biological sciences, specializing in biochemistry, 2008, Institute of Biochemistry and Biophysics PAS). Professional experience: 2008–2010 Assistant at the Institute of Biochemistry and Biophysics, since September 2011, Assistant Professor in the Department of Drug Technology and Biotechnology. Awards/prizes: Award of the Scientific Council of IBB PAS for the laureate of the FNP POMOST program in 2013. Number of publications: 12 publications in journals from the philadelphia list. Research interests: biology of pathogenic bacteria and yeast, the use of yeast in the production of selected primary and secondary metabolites.

e-mail: jmierzejewska@ch.pw.edu.pl, phone: +48 22 234 7074